

REFERENCES AND FURTHER READING

- **NICE** OHT diagnosis and management: NICE Guideline NG81 (2017). Glaucoma: diagnosis and management <https://www.nice.org.uk/guidance/ng81>
- **SIGN** Scottish Intercollegiate Guidelines Network (SIGN). Glaucoma referral and safe discharge. Edinburgh: SIGN; 2015. (SIGN publication no. 144, March 2015) <http://www.sign.ac.uk>
- **CMG's** Clinical Management Guidelines College Optometrists Glaucoma <https://www.college-optometrists.org/guidance/clinical-management-guidelines.html>
- **OHTS** Kass MA, Heuer DK, Higginbotham EJ, et al. The Ocular Hypertension Treatment Study: a randomized trial determines that topical ocular hypotensive medication delays or prevents the onset of primary open-angle glaucoma. *Arch Ophthalmol.* 2002;120:701-713.
- **LIGHT** Gazzard G, et al. Laser in Glaucoma and Ocular Hypertension (LiGHT) trial. A multicentre, randomised controlled trial: design and methodology. *Br J Ophthalmol.* 2018 May;102(5):593-598
- Nelson P, Aspinall P, O'Brien C Patients' perception of visual impairment in glaucoma: a pilot study *British Journal of Ophthalmology* 1999;83:546-552
- Leske MC, Connell AM, Wu SY, et al. Risk factors for open-angle glaucoma. The Barbados Eye Study. *Archives of Ophthalmology* 1995;113: 918-2
- de Voogd S, Ikram MK, Wolfs RC, et al. Incidence of open-angle glaucoma in a general elderly population: the Rotterdam Study. *Ophthalmology* 2005; 112: 1487-93
- Klein BE, Klein R, Sponsel WE, et al. Prevalence of glaucoma. The Beaver Dam Eye Study. *Ophthalmology* 1992; 99: 1499-1504
- Mitchell P, Smith W, Attebo K, Healey PR. Prevalence of open-angle glaucoma in Australia. The Blue Mountains Eye Study. *Ophthalmology* 1996; 103: 1661-9