

References

1. Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, Wong TY, Naduvilath TJ, Resnikoff S. Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050. *Ophthalmology*. 2016 May 1;123(5):1036-42.
2. Pärssinen O, Soh ZD, Tan CS, Lanca C, Kauppinen M, Saw SM. Comparison of myopic progression in Finnish and Singaporean children. *Acta Ophthalmologica*. 2021 Mar;99(2):171-80.
3. Németh J, Tapasztó B, Aclimandos WA, Kestelyn P, Jonas JB, De Faber JT, Januleviciene I, Grzybowski A, Nagy ZZ, Pärssinen O, Guggenheim JA. Update and guidance on management of myopia. European Society of Ophthalmology in cooperation with International Myopia Institute. *European Journal of Ophthalmology*. 2021 May;31(3):853-83.
4. Lam CS, Tang WC, Lee PH, Zhang HY, Qi H, Hasegawa K, To CH. Myopia control effect of defocus incorporated multiple segments (DIMS) spectacle lens in Chinese children: results of a 3-year follow-up study. *British Journal of Ophthalmology*. 2022 Aug 1;106(8):1110-4.
5. Chamberlain P, Bradley A, Arumugam B, Hammond D, McNally J, Logan NS, Jones D, Ngo C, Peixoto-de-Matos SC, Hunt C, Young G. Long-term effect of dual-focus contact lenses on myopia progression in children: A 6-year multicenter clinical trial. *Optometry and Vision Science*. 2022 Mar 1;99(3):204-12.
6. Si JK, Tang K, Bi HS, Guo DD, Guo JG, Wang XR. Orthokeratology for myopia control: a meta-analysis. *Optometry and Vision Science*. 2015 Mar 1;92(3):252-7.
7. Gordon RA, Donzis PB. Refractive development of the human eye. *Archives of ophthalmology*. 1985 Jun 1;103(6):785-9.
8. Ip JM, Huynh SC, Kifley A, Rose KA, Morgan IG, Varma R, Mitchell P. Variation of the contribution from axial length and other oculometric parameters to refraction by age and ethnicity. *Investigative ophthalmology & visual science*. 2007 Oct 1;48(10):4846-53.
9. McCullough S, Adamson G, Breslin KM, McClelland JF, Doyle L, Saunders KJ. Axial growth and refractive change in white European children and young adults: predictive factors for myopia. *Scientific reports*. 2020 Sep 16;10(1):15189.
10. Morgan PB, McCullough SJ, Saunders KJ. Estimation of ocular axial length from conventional optometric measures. *Contact Lens and Anterior Eye*. 2020 Feb 1;43(1):18-20.

References

11. Tideman JW, Snabel MC, Tedja MS, Van Rijn GA, Wong KT, Kuijpers RW, Vingerling JR, Hofman A, Buitendijk GH, Keunen JE, Boon CJ. Association of axial length with risk of uncorrectable visual impairment for Europeans with myopia. *JAMA ophthalmology*. 2016 Dec 1;134(12):1355-63.
12. COMET Group. Myopia stabilization and associated factors among participants in the Correction of Myopia Evaluation Trial (COMET). *Investigative ophthalmology & visual science*. 2013 Dec;54(13):7871.
13. Bullimore MA, Lee SS, Schmid KL, Rozema JJ, Leveziel N, Mallen EA, Jacobsen N, Iribarren R, Verkicharla PK, Polling JR, Chamberlain P. IMI—onset and progression of myopia in young adults. *Investigative Ophthalmology & Visual Science*. 2023 May 1;64(6):2-.
14. Khan HA, Naduvilath T, Tahhan N, Sankaridurg PR. Myopia Progression in Adults: A Retrospective Analysis. *Optometry and Vision Science*. 2023 Aug 1;100(8):537-42.
15. Jorge J, Almeida JB, Parafita MA. Refractive, biometric and topographic changes among Portuguese university science students: a 3-year longitudinal study. *Ophthalmic and Physiological Optics*. 2007 May;27(3):287-94.
16. Morgan IG, Wu PC, Ostrin LA, Tideman JW, Yam JC, Lan W, Baraas RC, He X, Sankaridurg P, Saw SM, French AN. IMI risk factors for myopia. *Investigative ophthalmology & visual science*. 2021 Apr 28;62(5):3-.
17. Lee SS, Mackey DA. Prevalence and risk factors of myopia in young adults: review of findings from the raine study. *Frontiers in Public Health*. 2022 Apr 27;10:861044.
18. Sha J, Tilia D, Diec J, Fedtke C, Yeotikar N, Jong M, Thomas V, Bakaraju RC. Visual performance of myopia control soft contact lenses in non-presbyopic myopes. *Clinical optometry*. 2018 Jul 25:75-86.
19. Vestergaard AH, Hjortdal JØ, Ivarsen A, Work K, Grauslund J, Sjølie AK. Long-term outcomes of photorefractive keratectomy for low to high myopia: 13 to 19 years of follow-up. *Journal of Refractive Surgery*. 2013 May 1;29(5):312-9.